

Inactivated H1N1 Vaccine

Vaccine Description	<ul style="list-style-type: none"> • Monovalent H1N1 vaccine; uses same process as inactivated influenza vaccine • Contains egg protein. See package insert. 	
Dose & Route	<ul style="list-style-type: none"> • Dose for age 3 years and older: 0.5 mL • Route for all doses: IM (Precaution: hemophilia, thrombocytopenia, and anticoagulation therapy) 	
Indications	<p>Initial Target Groups:</p> <ul style="list-style-type: none"> • Pregnant women • Persons who live with or provide care for infants aged < 6 months • Healthcare and emergency medical services personnel • Persons 6 months through 24 years • Persons 25 through 64 years of age who have medical conditions that put them at risk for influenza-related complications. <p>As supply improves, expand to all persons 25 through 64 years of age, followed by persons 65 years of age and older.</p>	
Administration Schedule	Dose	Recommended Interval
Adults	One dose: 0.5 mL	Single dose
Contraindications	<ul style="list-style-type: none"> • Serious allergic reaction to prior dose, influenza vaccine, vaccine component, or to eggs • Moderate or severe acute illness • Prior serious adverse event or history of Guillain-Barré syndrome (GBS) within 6 weeks of a previous dose of influenza vaccine 	
VIS: not available yet		

Live Attenuated H1N1 Vaccine

Vaccine Description	<ul style="list-style-type: none"> • Live attenuated, monovalent, nasally administered vaccine; uses same processes as LAIV • Contains egg protein. See package insert. 	
Dose & Route	<ul style="list-style-type: none"> • Dose: 0.2 mL • Route: Intranasal (half per nostril) 	
Indications	<p>Initial Target Groups:</p> <ul style="list-style-type: none"> • People who live with or provide care for infants aged < 6 months who do not have a contraindication • Healthcare and emergency medical services personnel who do not have a contraindication • People 2 through 24 years who do not have a contraindication <p>As supply improves, expand to all persons 25 through 49 who do not have a contraindication</p>	
Administration Schedule	Dose	Recommended Interval
Adults	0.2 mL	Single dose
Contraindications	<ul style="list-style-type: none"> • History of hypersensitivity, especially anaphylactic reactions, to prior dose or any component, including eggs or egg products • Moderate or severe acute illness • History of Guillain-Barré syndrome within 6 weeks of influenza vaccine • People who are immunosuppressed due to disease or medical treatment • Pregnancy • People with chronic conditions that place them at high risk for complications from influenza illness (e.g., heart disease, pulmonary disease, diabetes, renal disease, sickle cell anemia) 	
Special Considerations	<ul style="list-style-type: none"> • Defer administration if nasal congestion might prevent vaccine from reaching nasopharyngeal mucosa • May be given at the same time as other live vaccines, except LAIV. • See Storage and Handling Section 	
VIS: Not available yet		