

Inactivated H1N1 Vaccine

Vaccine Description	<ul style="list-style-type: none"> • Brands: Vaccine from sanofi pasteur approved for 6 months and older; Vaccine from Novartis approved for 4 years of age and older • Monovalent H1N1 vaccine; ; uses same process as inactivated influenza vaccine • Contains egg protein. See package insert. 	
Dose & Route	<ul style="list-style-type: none"> • Dose for age 6 months to 35 months: 0.25 mL • Dose for age 3 years and older: 0.5 mL • Route for all doses: IM (Precaution: hemophilia, thrombocytopenia, and anticoagulation therapy) 	
Indications	<ul style="list-style-type: none"> • All children from 6 months through 18 years of age <p>If very limited vaccine supply:</p> <ul style="list-style-type: none"> • All children 6 months through 4 year of age • All children 5 through 18 years of age who have medical conditions that put them at higher risk for influenza-related complications 	
Administration Schedule	Dose	Recommended Interval
6 months through 9 years of age	6 to 35 months: 0.25 mL Older than 3 years: 0.5 mL	Give 2 doses separated by about 4 weeks
10 years of age and older	One dose: 0.5 mL	Single dose
Contraindications	<ul style="list-style-type: none"> • Serious allergic reaction to prior dose, influenza vaccine, vaccine component, or to eggs • Moderate or severe acute illness • Prior serious adverse event or history of Guillain-Barré syndrome (GBS) within 6 weeks of a previous dose of influenza vaccine 	
VIS: Not available yet		

Live Attenuated H1N1 Vaccine

Vaccine Description	<ul style="list-style-type: none"> • Brand: FluMist® • Live, monovalent, nasally administered vaccine; uses same processes as LAIV • Contains egg protein. See package insert. 	
Dose & Route	<ul style="list-style-type: none"> • Dose: 0.2 mL Route: Intranasal (half per nostril) 	
Indications	<ul style="list-style-type: none"> • Healthy children from 2 through 18 years of age <p>If very limited supply of vaccine: Healthy children 2 through 4 year of age</p>	
Administration Schedule	Dose	Recommended Interval
2 through 9 years of age	0.2 mL	Give 2 doses separated by about 4 weeks
10 years of age and older	0.2 mL	Single dose
Contraindications	<ul style="list-style-type: none"> • History of hypersensitivity, especially anaphylactic reactions, to prior dose or any component, including eggs or egg products • Children and adolescents (2 through 18 years of age) receiving chronic aspirin or salicylate-containing medication therapy because of the risk for Reye syndrome • Moderate or severe acute illness • History of Guillain-Barré syndrome within 6 weeks of influenza vaccine • People who are immunosuppressed due to disease or medical treatment • Pregnancy • People who have asthma, reactive airway disease, or other chronic pulmonary disease • People with chronic conditions that place them at high risk for complications from influenza illness (e.g., heart disease, diabetes, renal disease, sickle cell anemia) 	
Special Considerations	<ul style="list-style-type: none"> • Defer administration if nasal congestion might prevent vaccine from reaching nasopharyngeal mucosa • May be given at the same time as other live vaccines, except LAIV. • See Storage and Handling Section 	
VIS: Not available yet		

NEW